

Fighting Fatigue:

A Guide for

Cardiothoracic Surgery Residents



Thoracic Surgery Directors Association

Thoracic Surgery Residents Association

**Thoracic Surgery Directors Association
Thoracic Surgery Residents Association
633 N. St. Clair Street, 23rd Floor
Chicago, IL 60611
312-202-5854
773-289-0871 (fax)
tsda@tsda.org
www.tsda.org
www.tsrnet.org**

Adapted from: Royal College of Physicians. *Working the night shift: preparation, survival and recovery. A guide for junior doctors.* London: Royal College of Physicians, 2006.

Copyright © 2006 Royal College of Physicians. *Adapted by permission.*

Prepared on behalf of a multidisciplinary Working Group by Nicholas Horrocks MSc and Roy Pounder MD DSc FRCP.

Edited by Errol Bush, MD (University of California San Francisco) and Stephanie Mick, MD (Brigham and Women's Hospital).

Contents

Key points	4	
Introduction	5	
Sleep and shift work		5
Night work and safety		6
Preparing for a night call rotation	7	
Successful sleep at home		7
Getting plenty of sleep before a night call rotation begins		7
Taking an afternoon nap		8
Can you adapt to night work?		8
Surviving the night shift	9	
Maintaining your alertness and vigilance while on duty		9
Napping while on duty		9
Bright light		10
Eating at night		10
Caffeine		10
Recovering from the night shift	12	
Getting home from work		12
Working further night shifts		12
Before you go to bed		13
Sleeping in the daytime		13
Recovering after your final night shift		14
The way ahead	14	
References	15	
Members of the Working Group	17	

KEY POINTS

Maximizing your ability to cope with night rotations

Facts

- Night shifts require doctors to be alert when their bodies tell them to sleep.
- Working at night generates an increasing sleep debt.
- Sleep-deprived residents have more attentional failures and make more clinical errors and incorrect diagnoses.
- Residents have more road traffic accidents when tired.
- Exhaustion erases recent learning.

Key advice

Preparing for the night shift

- Build a successful normal sleep routine.
- Get extra sleep before working the first night shift.
- Take a two-hour afternoon nap before coming on duty.

Surviving the night shift

- Take 20- to 45-minute naps to counteract fatigue.
- Your alertness will be improved by exposure to bright light during the night.
- Do not miss routine meals when working at night.
- Use caffeine cautiously, if at all, as it is a stimulant.

Recovering from the night shift

- If planning a long drive home, consider the risks.
- Once home, try to sleep immediately.
- Develop a routine for sleeping during the daytime.
- Keep your sleep debt to a minimum.

INTRODUCTION

Working at night is an essential part of providing a comprehensive 24-hour service to patients. However, night work requires residents to remain awake and alert when physiologically programmed to be asleep. Many residents may take 24-hour calls or primary night calls on certain rotations, and, on these occasions, residents can expect to stay away and working throughout the entire night.

The Accreditation Council for Graduate Medical Education (ACGME) Resident Duty Hours Standards were introduced to help protect patients and residents. This Guide does not deal with the appropriateness of the Resident Duty Hours Standards, rather it is designed to provide useful advice to residents working late at night.

• Night shifts require doctors to be alert when their bodies tell them to sleep.

Working at night can have consequences for both patient and personal safety, as it increases the risk of making poor decisions or even mistakes. It is therefore important to learn how to prepare for working at night and to manage your sleep, so that you minimize risk to yourself and to your patients.

Some of the suggestions in this Guide may be considered obvious but have been included for completeness. Others, such as the advice on napping, may be less obvious. However, they are all based on research, which is consistent and clear.

The aim of the Guide is to provide simple advice on how to prepare to work at night. It includes guidance on managing sleep at home, staying as alert and refreshed as possible while on duty at night, and how to recover from working nights in the most efficient and effective way. Although the Guide has been written with those working extended night or on-call rotations in mind, much of the advice is applicable to those residents taking more prolonged calls. Indeed, these longer calls carry the risk of even more severe sleep deprivation, than simple night call.

Sleep and shift work

Many people are expected to work at night, and most do so relatively successfully. However, all have to cope with the fact that working at night inevitably causes sleep deprivation and fatigue. This is because the human body is designed to sleep at night.

Our bodies are controlled by an internal daily body clock, situated in the suprachiasmatic nucleus (SCN) in the hypothalamus. The SCN spontaneously generates the circadian rhythms that regulate many physiological and behavioral processes in our bodies, such as temperature control, hormone production, alertness and sleep.

These circadian rhythms run over a period of approximately 24 hours and are strongly influenced by the natural cycles of light and dark. At night, many of the processes that are active during the day start to slow down as our bodies prepare for sleep. The circadian pacemaker also stimulates night-time release of the 'sleep hormone' melatonin from the pineal gland, which lowers alertness and increases the desire for sleep.

Working at night involves fighting against these rhythms, and trying to be alert when you are programmed to be asleep. In addition, when a night call finishes and you go home to try and sleep, the cues from your internal body clock, daylight, and society in general, all tell you that it is time to be active. Your sleep is likely to be fragmented and brief, no matter how tired you may feel. Crucially, you will not make up all of the hours of sleep that you have lost during the previous night.

Although individuals vary, most people need at least one hour of sleep for every two hours awake, or approximately eight to nine hours of sleep each night. If you manage less than this (as occurs in shift workers, who have sleep of poorer quality and shorter duration than non-shift workers¹) then you will incur a 'sleep debt'. This sleep debt is cumulative, so the more sleep you miss, the greater in debt you will be. The only way to repay the debt is by catching up on the lost sleep, and the sleep debt must be repaid soon after it is incurred. A fundamental aspect of being a successful night worker is learning how to manage your daytime sleep (and fatigue at night) so that you keep your sleep debt to a minimum.

• **Working at night generates an increasing sleep debt.**

Night work and safety

Fatigue has long been known to reduce performance. If you work at night you are trying to function when your alertness, vigilance, and cognitive reasoning are at their lowest. This applies to residents just as much as it does to other night workers.

In fact, there is now clear evidence that residents who are sleep-deprived have more attentional failures and make more clinical errors than when they are able to gain enough sleep.²⁻⁶ Furthermore, 20-25 hours without sleep – as might be experienced by a resident who has just worked 24-hour call – reduces psychomotor performance to the level of someone with a blood alcohol concentration of 0.10%.⁷⁻¹⁰ This is even greater than the level (0.08%) at which it becomes illegal to operate a motor vehicle in the United States.

• **Sleep-deprived residents have more attentional failures, and make more clinical errors and incorrect diagnoses.**

Evidence has shown that residents who take call 24 hours or longer more than double their risk of being involved in a road traffic accident on their drive home compared with those taking shorter calls.¹¹ The likelihood of crashing on the way home is also greater following a night shift than after other shifts.^{12,13} Moreover, when you are tired you become less able to judge your own performance accurately, so you may not even realize that you are making mistakes.¹⁴

• **Interns have more road traffic accidents when tired.**

Exhaustion also impairs recent learning^{15,16} and has been shown to decrease the ability of residents to make correct diagnoses,³ with important implications for both training and medical care in general.

• **Exhaustion erases recent learning.**

PREPARING FOR A NIGHT CALL ROTATION

The combination of fatigue and a poorly adapted body clock makes working during the night uncomfortable and increases errors. Preparing to work overnight is all about making sure you are rested and refreshed as possible before coming on duty.

Successful sleep at home

An essential first step is to manage your normal sleep when at home. Whether you are on a night call rotation or not, make sure that your bedroom is a suitable place in which to sleep.

- **Build a successful normal sleep routine.**

Most importantly, try to associate your bedroom with sleeping. Avoid watching the television, using a computer or smartphone, or playing videogames in the bedroom. Whenever you try to sleep at home, the bedroom is where you should go, rather than curling up on the sofa or in a chair. You will sleep best lying down in bed.

Although there may be matters that demand your attention, medical or otherwise, when you are trying to fall asleep, it is helpful to try consciously not to worry. Try not to let your mind dwell on the upcoming or previous call. If possible, you should actively put worrying concerns out of your mind and concentrate on pleasant thoughts, or focus your mind on an innocuous but absorbing activity such as mentally walking a route through a favorite park.

If you cannot sleep after having been in bed for 30 minutes or so, get up and go to another room and do something to distract yourself. Try some relaxation exercises, listen to soothing music, or perhaps take a bath. When you feel tired again, get back into bed and try to sleep again. Do not lie in bed stressed about the fact that you cannot sleep – this will not help.

It is important to try and build positive associations between being in bed and sleeping. If you can do this, your ability to fall asleep once you do get into bed will be improved.

Getting plenty of sleep before a night call rotation begins

Once you have established a successful sleep routine, make the most of it. Many people fail to get enough sleep, both before a call rotation and in general, so try to make sure that you are as well rested as you can be before the night call rotation begins. Any sleep that you have missed before you start is unlikely to be made up during the time that you are working at night. The greater your sleep debt, the more fatigued you will be, and the worse you are going to feel.

- **Get extra sleep before working at night.**

Remember, if you have not slept or rested at all since waking the previous day, by the time you come off the first night call you may well have been awake for 24 hours or more. To avoid this, try to sleep until at least midday on the morning before you take the first call if possible. Some people also stay up later the previous evening in order to begin to adapt their body clock and to make sleeping later easier. However, keep in mind that a late night out drinking will make you sleep poorly and will tend to increase your sleep debt and fatigue the next day.

Taking an afternoon nap

In addition to sleeping late, taking an afternoon nap is an extremely important way of making sure you are well rested before you start a night call. A pre-call two-hour nap will reduce the build-up of fatigue and make it much easier to remain awake and functional during the middle of the night.

Take your nap in the afternoon rather than just before coming on for a night call, because early evening is one of the times when your body is most alert and so sleep will be more difficult. By resting in the late afternoon, you can take advantage of the fatigue that you have already built up to help you sleep and to maximize your alertness throughout the night. Ideally, this rest should last at least two hours, to incorporate a beneficial period of deep sleep.

- **Take a two-hour afternoon sleep before coming on duty.**

Can you adapt to night work?

Body clock adjustment is very unlikely to occur in residents on short night call rotations. Given this, perhaps the most important thing to remember is that you need to take an active approach to managing sleep and fatigue. This is particularly true for the lifestyle of residents. As a resident, you will get tired, and you will become sleep-deprived, especially if you work several consecutive nights in a row. Inevitably, this will affect both how you feel and how you perform.

However, by preparing yourself sufficiently in advance, mentally and physically, you can reduce the negative impact of working at night on your well-being. This will not only make the experience less painful (or even enjoyable), but also safer for you, your patients, and those around you.

SURVIVING THE NIGHT SHIFT

Actually staying awake when you are in the hospital will depend on how much work you have to do. However, your levels of alertness and vigilance will be much lower than normal, so maintaining your performance at a safe level should be your priority.

Maintaining your alertness and vigilance while on duty

The natural circadian nadir is in the middle of the night, between 3 am and 6 am. During this period, the body is programmed to be at its least active, and workload in the hospital also tends to be at its lowest. The lighter activity, especially at this time, may make it more difficult to stay awake, and so this period of the night may well be when you feel most inclined to sleep.

Napping while on duty

Developing a napping routine is an indispensable part of working safely overnight. A ‘prophylactic’ afternoon nap before you come on duty will help keep minimize fatigue, but taking a nap during the night is essential for maintaining vigilance and alertness. Naps are powerful means of staying refreshed,^{17,18} both before and while on duty, and even naps as short as 20 to 45 minutes have been shown to provide positive benefits.¹⁹

• Take 20- to 45-minute naps to counteract fatigue.

Your nightly naps should last no longer than 45 minutes. This is because there are different stages of sleep, with cycles of 90- to 100-minutes’ duration. Set an alarm before you nap to make sure you do not fall into a prolonged deep sleep, and, if possible, give yourself enough time to recover prior to resuming your duties. By limiting the length of your nap, it is possible to avoid awakening during a period of deep sleep, when the general effort of awakening and any associated sleep inertia will be much greater. This is important for a resident who may need to be fully alert without warning.

Naps are most effective if taken early, before you feel really tired, and should be taken in surroundings that are quiet and dark. Try to lie down, or have your legs supported. It will be much harder to rest if you cannot at least recline.²⁰

If you are working a series of consecutive night shifts, try to avoid prolonged sleeping (rather than just napping) during the night. The more sleep that you get at night, the harder it will be to do so during the day when you have the opportunity for longer cycles of potentially unbroken and beneficial deep sleep. If you have periods of inactivity during a night shift you should offer help to other colleagues; repeated inactivity should lead to a review of rostering arrangements.

Bright light

Plan to maximize your exposure to light throughout the night shift. Exposure to light during the night, including indoor light from a bright desk lamp or normal overhead lights, has an alerting effect on the brain and improves performance.²¹

- **Your alertness will be improved by exposure to bright light during the night.**

Other than napping, try to make your medical area brightly illuminated, especially when working. Importantly, intermittent light exposure is nearly as effective as continual exposure.²² Even if you can only be exposed to indoor light from time to time through the shift, it will still be beneficial.

Eating at night

Eat and drink properly so that you do not start your night shift hungry or dehydrated. It is very easy when working at night to miss routine meals, because circadian patterns affect appetite, and dining facilities are often limited or closed. Ideally, you should try to maintain a similar eating pattern to the one you follow during the day. There is some evidence that a high-protein, low-carbohydrate meal is best for maintaining the night shift alertness.²³

- **Do not miss routine meals when working at night.**

Eat a full meal before you come on duty, have 'lunch' halfway through your shift, and finally enjoy an easily digestible meal before trying to sleep when you are at home, if you feel hungry then. If your hospital does not have adequate facilities for providing good meals at night, then bring in your own food.

Caffeine

Some residents use caffeine as a stimulant to help them to stay awake. Despite its widespread use, caffeine does have side effects and repetitive use is discouraged. Depending on your tolerance, too much caffeine can cause gastrointestinal upsets and muscle shakes. Additionally, it should not be taken at least four hours before the end of a night shift, since its long-lasting effects may cause you to find it harder to sleep once you get home.

If you do decide to use caffeine to aid your alertness, it may be best to take it in small amounts.²⁴ The effects of a cup of coffee can start being felt within as little as 20 minutes, and may last for up to three or four hours, depending on the individual and the type of coffee. Likewise, caffeine-containing energy drinks may help you to stay alert. By taking an appropriate small dose of caffeine just before you nap, its effects should start to be felt about the time that you return to duty. The caffeine may also help to overcome the transient sleep inertia you feel after the nap.

- **Use caffeine cautiously, if at all, as it is a stimulant.**

The caffeine content of some common drinks and food.²⁵

Drink/food	Caffeine content (mg)
Average cup of instant coffee (200 ml)	75
Average cup of brewed coffee (200 ml)	100-250
Average cup of tea (200 ml)	50
Herbal tea	0
Decaffeinated tea or coffee	3-5
Hot chocolate	5-7
Coca-Cola (330 ml can)	32
Diet Coke (330 ml can)	42
Pepsi (330 ml can)	35
Diet Pepsi (330 ml can)	34
Red Bull (250 ml can)	80
Bar of dark chocolate (50 g)	up to 50
Bar of milk chocolate (50 g)	up to 25
Pro Plus caffeine tablets	50 mg per tablet

RECOVERING FROM THE NIGHT SHIFT

Your actions at the end of a night shift should depend on whether or not it is your last shift. If you are working further night shifts, then it is important to focus on keeping your sleep debt under control. However, if your night duties have finished you should aim to repay any sleep debt you have built up, before getting back to your normal daytime life.

Getting home from work

Once your shift is over, it is likely that your first thought will be to get out of the hospital and go home. If you are planning to drive a long distance, just consider whether it is wise. Exhausted drivers kill both themselves and others, and driving fatigued is effectively no different than driving while over the legal limit for blood alcohol concentration.⁷⁻⁹ If you have worked overnight you will have slowed reflexes and poor judgment, or you may fall asleep at the wheel. Once you finish your shift you are likely to become exhausted very quickly and the longer your journey home, the greater the risk that you will cause an accident.²⁶⁻²⁷

When you are tired, your ability to judge your own performance is impaired, so you may think that you are better able to cope with driving than you actually are. In fact, the more tired you are, the less able you are to accurately assess your own performance.¹⁴ Furthermore, you cannot judge the exact point when you will fall asleep even though you can anticipate broadly when it is going to happen – it is time to pull off the road if you are fighting to keep your eyes open. Despite the inconvenience and cost, it may be better to either use public transportation to ensure that you arrive home safely, or to use sleep accommodations. Recently, some programs have begun to offer complimentary taxi vouchers for fatigued residents to their home, but you should check with your program coordinator in advance to see which specific options are available for your program.

• If planning a long drive home, consider the risks.

Working further night shifts

If you have to work more nights and are not driving, wear dark sunglasses on your way home to minimize your exposure to sunlight.²¹ Bright light is one of the key triggers for resetting your internal body clock back to its normal daytime schedule, and it should be avoided if you need to sleep.

As soon as you get home the best thing to do is try and sleep. Your aim should be to fully recuperate as soon as possible before your next shift, and to keep your body on a night work setting.

Before you go to bed

When you get home, don't get distracted by other things that cause you to delay going to bed. The longer that you delay, the more awake you are likely to become and the harder you will find it to sleep, no matter how tired you may be. Shift workers who go to bed at 10 am tend to sleep for at least four hours, whereas those who go to sleep at midday sleep for an hour less.²⁸

• **On getting home, try to sleep immediately.**

If you are hungry or thirsty, however, have something to eat and drink. You don't want to be awoken from precious daytime sleep by feelings of hunger or thirst.

Avoid alcohol, because although its relaxing effects may help you to fall asleep initially, the quality of your sleep will be diminished and you may well suffer from insomnia. Alcohol disturbs the stages of deep sleep. When you wake up you will not feel refreshed and, more significantly, your sleep debt will not have been greatly reduced.

If you are a smoker then it is also a good idea to avoid smoking before you try to sleep. Nicotine is a stimulant and it will make it more difficult for you to sleep. Avoid any activities that may increase your alertness until the hours before your next shift.

Sleeping in the daytime

Your bedroom should be quiet, dark, and cool. Noise, bright sunlight, and temperature are common complaints of people unable to sleep during the day. Make sure you have good blackout curtains that filter out all external light, or use eyeshades. Wear expandable foam earplugs if necessary to block out daytime noises like traffic and building work. Keep the room cool; an electric fan will not only circulate air, but can also provide a neutral and constant background noise. A soothing CD playing at very low volume may be helpful.

Turn off your mobile phone, disconnect any landline or switch on voicemail. If it is safe to do so, consider putting a notice on your front door to warn others that you are sleeping.

• **Develop a routine for sleeping during the daytime.**

Sleeping tablets are not recommended to keep you asleep after a night shift, because of their potential hangover and addictive effects. Consult your general practitioner if you feel that they are absolutely necessary, but certainly do not self-prescribe. Sleeping in the daytime is more difficult than sleeping at night, but many people manage it successfully. It is far better to review your sleep routine and sleeping arrangements than to rely on sedation.

If you wake up earlier than intended, just relax and you may fall asleep again. Otherwise, get up and take it easy. Remember to have at least a two-hour sleep before going back on duty, and make sure you eat and hydrate properly. The most important thing is to try and keep your sleep debt to a minimum, so the more daytime sleep that you get, the better you are going to feel. Reserve your recreation and stimulation for the early evening, and only expose yourself to bright light once you are ready to go to work.

• **Keep your sleep debt to a minimum.**

Recovering after your final night shift

If this is your final night shift, remember that the more consecutive nights you have worked, the greater your cumulative sleep debt is likely to be. Reducing the build-up of sleep debt and repaying this sleep debt promptly will help you to recover earlier, and may also have longer-term health benefits.

Sleep when you get home from work to get over some of your initial fatigue. When you wake up, get out of bed and do normal daytime things. Make sure you receive some exposure to daylight, but go to bed early that night and use this time to start really catching up on sleep. If you can sleep in the next morning then do so, but then get up and get on with your life. An early night on the following evening will help you catch up on more missed sleep, but the sooner you get back into your daily 'daytime' routine, the sooner your sleep patterns will return to normal.

THE WAY AHEAD

Few residents look forward to working overnight. Nevertheless, because healthcare is a 24-hour service in almost every specialty, a proportion of residents will always need to care for patients at night, and the experience can be very rewarding. Whether these hours are worked as night shifts or on-call, the risks associated with working during the biological night remain, and need to be approached, systematically. Each person is different, so finding the best combination of techniques for you may require some time. We hope the advice in this Guide will make the challenge of these duties not only a bit easier to tolerate, but also safer for both hospital patients and you, their doctor.

REFERENCES

1. Knauth P, Rutenfranz J. Duration of sleep related to the type of shiftwork. In: Reinberg A, Vieux N, Andlauer P (eds), *Advances in the Biosciences, Vol 30. Night and shiftwork: biological and social aspects*. New York: Pergamon Press, 1980: 161-8.
2. Lockley SW, Cronin JW, Evans EE, Cade BE *et al*. Effect of reducing interns' weekly work hours on sleep and attentional failures. *N Eng J Med* 2004; 351:1829-37.
3. Landrigan CP, Rothschild JM, Cronin JW, Kaushal R *et al*. Effect of reducing interns' work hours on serious medical errors among interns in intensive care units. *N Eng J Med* 2004;351:1838-48.
4. Friedman RC, Bigger JT, Kornfeld DS. The intern and sleep loss. *N Eng J Med* 1971; 285:201-3.
5. Grantcharov TP, Bardram L, Funch-Jensen P, Rosenberg J. Laproscopic performance after one night on-call in a surgical department: prospective study. *BMJ* 2001;323:1222-3.
6. Eastridge BJ, Hamilton EC, O'Keefe GE, Rege RV *et al*. Effect of sleep deprivation on the performance of simulated laparoscopic surgical skill. *Am J Surg* 2003;186:169-74.
7. Dawson D, Reid K. Fatigue and alcohol performance impairment. *Nature* 1997; 388:235.
8. Lamond N, Dawson D. Quantifying the performance impairment associated with fatigue. *J Sleep Res* 1999;8:255-62.
9. Williamson AM, Feyer A-M. Moderate sleep deprivation produces impairments in cognitive and motor performance equivalent to legally prescribed levels of alcohol intoxication. *Occup Environ Med* 2000;57:649-55.
10. Arnedt JT, Owens J, Crouch M, Stahl J, Carskadon MA. Neurobehavioral performance of residents after heavy night call vs after alcohol ingestion. *JAMA* 2005;294:1025-33.
11. Barger LK, Cade BE, Ayas N, Cronin JW *et al*. Extended work shifts and the risk of motor vehicle crashes among interns. *N Eng J Med* 2005;352:125-34.
12. Steele MT, Ma OJ, Watson WA, Thomas HA Jr, Muelleman RL. The occupational risk of motor vehicle collisions for emergency medicine residents. *Acad Emerg Med* 1999;6:1050-53.
13. Akerstedt T, Peters B, Anund A, Kecklund G. Impaired alertness and performance driving home from the night shift: a driving simulator study. *J Sleep Res* 2005;14:17-20.
14. Van Dongen HP, Maislin G, Mullington JM, Dinges DF. The cumulative cost of additional wakefulness: dose-response effects on neurobehavioral functions and sleep physiology from chronic sleep restriction and total sleep deprivation. *Sleep* 2003;26:117-26.
15. Stickgold R, James L, Hobson JA. Visual discrimination learning requires sleep after training. *Nature Neurosci* 2000;3:1237-8.
16. Stickgold R. Sleep-dependent memory consolidation. *Nature* 2005;437:1272-78.
17. Dinges DF, Broughton RJ (eds). *Sleep and alertness: Chronobiological, behavioural, and medical aspects of napping*. New York: Raven Press, 1981:1-322.
18. Dinges DF, Orne MT, Whitehouse WG, Orne EC. Temporal placement of a nap for alertness: contribution of circadian phase and prior wakefulness. *Sleep* 1987;10:313-29.
19. Naithoh P. Minimum sleep to maintain performance: the search for sleep quantum in sustained operations. In: Stampi C (ed), *Why we nap*. Boston: Birkhäuser, 1992.
20. Cole RJ. Postural baroreflex stimuli may affect EEG arousal and sleep in humans. *J Appl Physiol* 1989;67:2369-75.
21. Yoon IY, Jeong DU, Kwon KB, Kang SB, Song BG. Bright light exposure and light attenuation in the morning improve adaptation of night shift workers. *Sleep* 2002;25:351-6.

22. Rimmer DW, Boivin DB, Shanahan TL, Kronauer RE *et al.* Dynamic resetting of the human circadian pacemaker by intermittent bright light. *Am J Physiol Regul Integr Comp Physiol* 2000;279:R1574-R1579.
23. Romon-Rousseau M, Lancry A, Poulet I, Frimat P, Furon D. Effect of protein and carbohydrate snacks on alertness during the night. In: Oginski A, Pokorski J, Rutenfranz J (eds), *Contemporary advances in shiftwork research*. Krakow: Medical Academy, 1987:133-41.
24. Wyatt JK, Cajochen C, Ritz-De Cecco A, Czeisler CA, Dijk DJ. Low-dosed repeated caffeine administration for circadian-phase-dependent performance degradation during extended wakefulness. *Sleep* 2004;27:374-81.
25. Food Standards Agency. *Advice for pregnant women on caffeine consumption*, 10 October 2001. www.food.gov.uk/news/pressreleases/2001/oct/caffeinepregnant
26. Philip P, Taillard MA, Quera-Salva B, Bioulac B, Akerstedt T. Simple reaction time, duration of driving and sleep deprivation in young versus old automobile drivers. *J Sleep Res* 1999;8:9-14.
27. Horne J, Reyner L. Vehicle accidents related to sleep: a review. *Occup Environ Med* 1999;56:289-94.
28. Folkard S. Circadian rhythms and shiftwork: adjustment or masking? In: Hekkens WThJM, Kierhok GA, Rietveld WJ (eds), *Trending in chronobiology*. Oxford: Pergamon Press, 1988.

MEMEBERS OF THE ROYAL COLLEGE OF PHYSICIANS WORKING GROUP

Speakers – Professor Simon Folkard, Université René Descartes, Paris; Dr. Christopher Landrigan, Brigham & Women’s Hospital & Harvard Medical School, Boston; Dr. Steven W Lockley, Brigham & Women’s Hospital & Harvard Medical School, Boston; Dr. Deborah Powell, New Zealand Resident Doctors Association; Dr. Barbara Stone, Centre for Human Sciences, QinetiQ.

Members and contributors to the Guide – Dr. Peter Alexander, Intensive Care Society Trainees Committee; Dr. Mary Armitage, Clinical Vice-President, Royal College of Physicians; Professor Michael Bannon, Postgraduate Dean, Oxford PGME; Mr. Mike Beattle, Senior Business Manager, Doctors Team, NHS Employers; Professor Dame Carol Black, President, Royal College of Physicians; Mr. Stuart Blackwell, Royal College of Physicians, Patient and Carers Forum; Dr. Declan Chard, Chair, Trainees Committee, Royal College of Physicians; Ms. Sarah Connelly, Head of Medical Workforce Development, North Central London SHA; Dr. Robert Coward, SpR Advisor, Royal College of Physicians; Professor Sir Alan Craft, President, Royal College of Paediatrics & Child Health; Chairman, Academy of Medical Royal Colleges; Dr. John Curran, Royal College of Anaesthetists; Mr. Roger Currie, Royal College of Surgeons (Edinburgh); Ms. Lindsey Dawson, Health and Community Care Research Unit; Ms. Sue Dean, Associate Director, NHS Workforce Projects; Mr. Simon Denegri, Director of Corporate Communications, Royal College of Physicians; Mr. Andrew Drakeley, Chair, Academy of Medical Royal Colleges Trainee Doctors Group; Professor Bill Dunlop, Chair, Joint Consultants Committee; Mr. Simon Eccles, National Clinical Lead for Hospital Doctors, NHS Connecting for Health; Mr. Martin Else, Chief Executive, Royal College of Physicians; Mr. Paul Evans, NHS patient; Dr. Sara Fairbairn, Trainee Committee, Royal College of Physicians; Dr. Anna Forrest-Hay, Faculty of Accident & Emergency Medicine; Dr. Ashley Fraser, Medical Deputy Chair, Trainees Committee, Royal College of Physicians; Dr. Richard Long, Councillor, Royal College of Physicians; Professor John Lowry, Royal College of Surgeons (England); Dr. Andrew Mitchell, Royal College of Paediatrics & Child Health; Ms. Alice Murray, Medical student, UCL; Dr. Mike Page, Consultant Physician, Royal Glamorgan Hospital; Dr. Amit Patel, Trainee Committee, Royal College of Physicians; Dr. Jane Pateman, Associate Dean, London Deanery; Dr. Dilip Raje, Royal College of Physicians, Patient and Carers Forum; Miss Wendy Reid, Postgraduate Dean, London Deanery; Dr. Mark Roberts, SpR in Respiratory Medicine, Derbyshire Royal Infirmary; Dr. Andrew Rowland, Deputy Chairman, UK Junior Doctor’s Committee, BMA; Mr. Dean Royles, Head of HR Capacity & Employment, NHS Workforce Directorate; Ms. Joan Russell, Safer Practice Lead, National Patient Safety Agency; Dr. John Scarpello, Deputy Medical Director, National Patient Safety Agency; Dr. Mark Simmonds, Specialist Registrar in Medicine, Queen’s Medical Centre, Nottingham; Mr. Dominic Slade, Secretary, Association of Surgeons in Training; Dr. David Snashall, President, Faculty of Occupational Medicine; Dr. David Stevenson, RAF Centre of Aviation Medicine; Mrs. Elaine Stevenson, Safer Practice Lead, National Patient Safety Agency; Dr. Avindan Veiraiah, Specialist Registrar in Clinical Pharmacology, Llandough Hospital; Dr. Chandi Vellodi, Councillor, Royal College of Physicians; Mr. Jim Wardrope, President Elect, Faculty of Accident & Emergency Medicine; Mr. Richard Warren, Honorary Secretary, Royal College of Obstetricians & Gynecologists; Mr. David Wells, WTD Lead, Maternity & Paediatrics, Care Services Improvement Partnership; Miss Melissa Whitten, Chair, Royal College of Obstetricians & Gynecologists, Trainees Committee; Dr. Martha Wyles, Chair, Trainees Committee, Royal College of Paediatrics & Child Health; Mr. Robert Wykes, Royal College of Physicians, Patient and Carers Forum; Dr. Sashidhar Yeluri, SHO Surgical Rotation, St James’ University Hospital, Leeds.